


MVTX Cost & Schedule and R&D

- ▶ A master cost & schedule project file being updated
 - LBNL - manpower and technical resources, fixed cost etc.
 - Will get similar updates from MIT and other institutions
 - Prepare for BNL review (physics + cost & schedule)
 - To update MVTX proposal later this year

- ▶ MAPS R&D @LANL
 - Received 5 MAPS chips + high-speed readout adaptors to build a telescope
 - Setting up “MOSAIC” high-speed readout test bench @LANL
 - Two RUV1 boards available ~ April/May, “CRU” w/ Altera evaluation boards
 - In process of obtaining FELIX from BNL

- ▶ MAPS Readout workfest scheduled @UT-Austin, 4/19-20
 - UT-Austin, LANL, LBNL, BNL experts

HF-topical group



- ▶ **HF in sPHENIX:** in particular B -meson and b -jets, provide differentiating sensitivity to collision VS radiative energy loss, access to HQ transport parameter of QGP, total cross section .
 - B -meson: access down to zero p_T , max sensitivity to HQ mass effect
 - b -Jet: less dependence on FF complication, probing parton energy and higher-scale
- ▶ **Topical group formed** in Apr 2016 focusing on establishing b -jet program. Many progresses thanks to diligent team of developers. **Delivered** first Geant4-based b -jet tagging performance and expanding into correlation studies.
- ▶ **In the new era of MVTX program**, aim to expand the program in HF-meson program as initiated by LBNL and LANL groups, and serve the detector consortium of MVTX
- ▶ Topical group wiki: https://wiki.bnl.gov/sPHENIX/index.php/Heavy_Flavor_Topical_Group

Brainstorming meeting, Mar 8

- ▶ <https://indico.bnl.gov/conferenceDisplay.py?confId=2932>
- ▶ About 20 sPHENIX collaborator and Ivan Vitev called in the discussion

Current development planning assumed two time-scale:

- Director's review – Apr/May
- DOE review – past summer

Brainstorming meeting for MVTX detector physics deliverable

chaired by Jin Huang (Brookhaven National Lab), Michael McCumber (Los Alamos National Laboratory), Xin Dong (Lawrence Berkeley National Laboratory), Ming Liu (Los Alamos)

Wednesday, March 8, 2017 from 13:30 to 14:55 (US/Eastern)
at Universe (2-160)

Description Meeting URL
<https://bluejeans.com/891109017>

To join via Phone:
1) Dial:
+1.408.740.7256
+1.888.240.2560(US Toll Free)
+1.408.317.9253(Alternate number)
(see all numbers - <http://bluejeans.com/numbers>)
2) Enter Conference ID: 891109017





Wednesday, March 8, 2017

13:30 - 13:45	MVTX Schedule and Strategy 15' Speaker: Dr. Ming Liu (Los Alamos) Material: Slides
13:50 - 14:05	HF-jet status and ideas 15' Speakers: Dr. Jin Huang (Brookhaven National Lab), Dr. Michael McCumber (Los Alamos National Laboratory) Material: Slides
14:10 - 14:25	HF-meson status and ideas 15' Speaker: Dr. Xin Dong (Lawrence Berkeley National Laboratory) Material: Slides
14:30 - 14:45	CMS HF measurement and implication to sPHENIX studies 15' Speaker: Yen-Jie Lee (Massachusetts Institute of Technology) Material: Slides
14:50 - 14:55	Simulation planning 5' Material: Wiki TODO list

Next few pages





Technical performances:

- Realistic implementation in Geant4
 - **In verification:** implement ladder structure in simulation – Tony F., Gaku M.
 - **By Apr (Done):** digitization of MAPS detector - Tony F.
 - **By end Apr:** Update tracking performance plots for MAPS, DCA and dp/p resolution - Tony F.
 - **By summer (?):** complete the pile-up simulation framework – Mike M., Yorito Y.
- *b*-jet tagging algorithm
 - **By summer, help needed :** Investigating full-detector fast simulation for *b*-jet simulation. Look into general packages e.g. [DELPHES](#).
 - **By summer:** Full calorimetry simulation with secondary vertexing tagger – Sanghoon L.
 - **By summer:** Full calorimetry simulation with high-DCA track counting – Haiwang Y.



Physics performances:

- ▶ Update non-prompt D meson performance projection
 - **Deliverable by end Apr**: update the Rcp and v2 plot with more realistic simulations for MB and peripheral collisions - Xin D., Xiaolong C.
- ▶ Explore complimentary B-hadron channels beyond non-prompt-D
 - **By summer**: Fast simulation for exclusive channels, B→J/Psi K, B→D pi - Xin D., Xiaolong C.
 - **By summer, help needed**: B→non-prompt J/Psi→e⁺/e⁻ and p+p triggering
- ▶ Inclusive b-jet R_{AA}
 - **By Apr (?)**: Update theory curve to RHIC energy – Cesar da S. contact Vitev group
 - **Deliverable by end Apr**: Update theory curve for RHIC energy for R_{AA} plot
- ▶ di-b-jet asymmetry
 - **By Apr**: Extract di-jet purity from Geant4 simulation - Haiwang Y.
 - **Deliverable by end Apr**: Apply di-jet purity to projection – Darren M., Haiwang Y.
- ▶ b-jet-non-prompt-D asymmetry:
 - **Deliverable by end Apr**: Produce uncertainty projection in fast simulation – Xuan L.





Additional topics help (always) wanted

- ▶ **HF-hadron chemistry:** e.g. high stat. Λ_c , to study HQ hadronization
- ▶ **HF-meson correlations**, e.g. D-D_{bar} azimuthal correlations, to enhance sensitivity to HQ-medium interaction; enhance M/pT ratio scale comparing to D-b-jet correlation.
- ▶ Explore **b-jet substructure tools**:
 - Exercise jet-grooming algorithm, FF. – in collaboration with Jet Structure group
 - Tagging gluon splitting via multi-decay vertex in inclusive b-jets.
- ▶ Explore **Charmed-quark jet**:
 - charm fragmentation, completes mass hierarchy. Select D meson formed late in formation
 - Try out prompt-D tagger (ALI-PREL-117896) and Corrected Secondary vertex (arXiv:1612.08972)
- ▶ Explore **tagged D-meson in correlation with opposite hard structure**
 - Tagging initial c-quark kinematics with correlations, including D-meson - jet correlation and D-meson - photon correlation
 - Study D-FF and formation of D-meson
- ▶ Further **b-jet tagging developments**
 - Try different strategy: Soft-lepton tagging
 - Optimize analysis methods: likelihood analysis and machine learning tool
- ▶ **Triggering** of B-mesons in p+p collisions
 - D→J/Psi →e⁺/e⁻, EMCal trigger. Explore work by Sasha L.
 - D meson calorimetry trigger, turn on.
 - Large DCA triggers?


Progresses since brain storming meeting – modeling of silicon trackers



Tony Frawley:
Approximation of MAPS diffusion
parametrization [arXiv:10053.3710]
Used in MAPS digitization,
Reproduced ALICE claim of 5 μm res.
<https://github.com/sPHENIX-Collaboration/coresoftware/pull/255>



(a) $h = 2 \mu\text{m}$




Gaku Mitsuka :

- Built INTT geometry model
- Improving estimation of material budget
- Debugging $r\phi$ cluster distribution

Progresses since brain storming meeting




Non-prompt D-meson correlations

- ▶ Xuan Li (LANL) also started investigation of correlation of b-jet in correlation of a non-prompt D_0 -meson ($\rightarrow \pi + K$)
- ▶ Goal: tag initial quark energy + vector for non-prompt D measurements, probe b-quark energy loss and fragmentation; access to lower z , cut-off comparing to di-b-jets correlation; help purity of b-quark-jet tagging
- ▶ Producing fast simulation results for pre-proposal



Updates in simulation meetings:





<https://indico.bnl.gov/conferenceDisplay.py?confId=2684>



Preliminary study

Path forward: summary


- ▶ Apr review/preproposal:
Solidifying results and studies presented in Feb-2017 pre-proposal
- ▶ Summer review/full proposal:
 - Expansion of selected topics: more realistic simulation, exclusive B-meson reconstruction
 - Addressing BNL-charged review comments
 - Many more HF capabilities need your help to develop
- ▶ Expect workfest around time of May collaboration meeting








Extra information



Delivered plots – HF-jet for Feb-2017 MAPS pre-proposal



Curve update request, not yet received



Observable Projections

B-jet tagging in p+p

B-jet tagging in 10%C Au+Au

HF Topical group organization

▶ Co-conveners

- Jin Huang (Brookhaven National Lab)
<jhuang@bnl.gov>
- Mike McCumber (Los Alamos National Lab)
<mccumber@bnl.gov>



▶ We are very fortunate to have a diligent team working on a wide spectrum of high-priority development

- More manpower are always welcomed and needed!

▶ Communication:

- Discussion email list:
<https://lists.bnl.gov/mailman/listinfo/sphenix-hf-jets-l>
- Wiki page under construction:
https://wiki.bnl.gov/SPHENIX/index.php/Heavy_Flavor_Topical_Group

▶ Meetings/Events




- Use weekly simulation meetings for updates, as many high-priority tasks involve software developments with tracking detector designs
<https://indico.bnl.gov/categoryDisplay.py?categId=88>
- Goal oriented irregular events:
 - MVTX brainstorming meeting, Mar 8
 - MAPS+HF-jet joint workfests, e.g. Jan 5-7 2017 @ Santa Fe
 - Pre-collaboration meeting work-fest on May 16-17, 2016
 - Initial TG meeting on Apr 22, 2016




Tagging b -jets in sPHENIX

Exploring three leading methods for sPHENIX b -jets identification and crosscheck


- ▶ Multiple large DCA tracks
- ▶ Secondary vertex and kinematic fits
- ▶ B -meson tagging via semi-leptonic decay or direct invariant mass reconstruction
 - Need volunteer



What affects performance curves



sPHENIX-proposal projection based on fast sim.
(already updated to Geant4 simulation)



Saturate efficiency at level determined by

- Control light decay-background
- Fake high DCA (mismatching, alignment) → realistic sim.

Performance turn-on at primary particle's DCA peak
Efficiency improve w/ higher DCA precision and tracking eff.

Initial b -jet fraction

Past activities:


b-jet tagging – High DCA track counting

► Short history


- Dennis and Haiwang implemented track counting tagger in the full Geant4 simulation
- Haiwang produced projection plot in Geant4 simulation.
- Systematically validating the Geant4-based track fit procedure, in order to optimize 3-D DCA and likelihood

► Next


- Reevaluate in HI background with HIJING embedding
- Optimizing cuts to suppress fake off-vertex tracks




Fast sim in sPHENIX Proposal



Full Geant4 Sim



Exploring 3-D DCA in G4



From Haiwang's talk

<https://indico.bnl.gov/conferenceDisplay.py?confId=1926>

b-tagging performance in HI


Past activities:

b-jet tagging – Secondary vertex

- ▶ Short history
 - Haiwang developed new Kalman filter (GenFit2) with vertex finder integration (RAVE)
 - Sanghoon implemented Secondary vertex finder in jet
 - $p+p$ performance plot used in tracking review
- ▶ Next:
 - Fixing a refitting inefficiency issue (further improve $p+p$ results)
 - Reevaluate in HI background with HIJING embedding




Secondary vertex *b*-tagger



From Sanghoon's talk


<https://indico.bnl.gov/conferenceDisplay.py?confId=1928>

Secondary vertex kinematics fits Data driven *b*-jet purity estimation





An vulnerability (opportunity) of HF-probes

sPHENIX scientific proposal






CMS, Phys.Rev.Lett. 113 (2014)
Phys.Lett. B726 (2013) 251-256



Lund String, Eur. Phys. J. C 17, 137–161 (2000)


b-quark jet selection: *b*-jet correlation

- ▶ Event topology to select *b*-quark jet
 - *b*-jet in correlation with opposite-going *B*-hadron, *b*-jet and photon
- ▶ sPHENIX provides good acceptance on *b*-di-jet and *b*-jet – non-prompt-*D* correlations
- ▶ Helps on purity of jet and *b*-tagging too
- ▶ Near term goals: fast-sim projections





b-jet + *B*-hadron, model

Physics Letters B750 (2015) 287–293




b di-jet, CMS 2016

CMS PAS HIN-16-005





di-jet acceptance in sPHENIX

sPHENIX scientific proposal




New studies for Di-b-jet asymmetry


With reference to recent CMS TN





CMS-HIN-16-005



Work started in Jan-2017 workfest
sPHENIX di-bjet asymmetry,
- Darren McGlinchey (UCB)



Di-*b*-jet asymmetry: sPHENIX projection



CMS-HIN-16-005, also Yen-Jie's talk
July 2016





On-going sPHENIX projection

- By Darren McGlinchey (UCB)
- Pythia8 (HardQCDBBar)
- Fast sim. (truth jets)
- Assuming di-*b*-jet tagging perf.
 - Efficiency 50%
 - High purity (100%)
- $R_{AA} = 0.6$ assumed
- sPHENIX proposal lumi. (100B MB)

● For $p + p$ use integrated luminosity of $\int \mathcal{L}_{pp} dt = 175 \text{ pb}^{-1}$

● For 0-10% Au+Au use $n + n$ equivalent luminosity of $\int \mathcal{L}_{nn} dt = N_{ev}^{AuAu} * \langle N_{coll} \rangle / \sigma_{nn} = 10\text{B} \times 962/42\text{mb} = 229 \text{ pb}^{-1}$





More ideas on b -quark jet selection?

Jet structure tools

- ▶ Jet structure tool developed in HEP adapted in HI field
- ▶ Jet grooming observable z_g to separate b -quark jet and $g/q \rightarrow b\bar{b}$ jet?
- ▶ **Mid-term goals:** in collaborate with JS TG in developing grooming tools – **volunteer welcomed!**


Earliest splitting:
More symmetric for $g/q \rightarrow b\bar{b}$ jet?



From Christof R.

<https://indico.bnl.gov/conferenceDisplay.py?confId=2683>




Alternate Detector Configurations



- Alternate detector configurations
 - Default -> 3 MAPS Layers, 4 ITT Layers, 40 Layers TPC
 - Light-> 2 MAPS Layers, 2 ITT Layers to save material budget
 - Slight performance advantage below 10GeV due to lower material budget
 - Minimal -> 1 Maps Layer, 2 ITT Layers, 40 Layers TPC
 - Significant performance decrease (remember 95% hit efficiency per layer)
 - No Maps
 - Likely death sentence for Heavy Flavor program...

Detector requirement on MAPS/MVTX

- ▶ Caveats: there are trade-offs between tail/efficiency/DCA. Important final check is b-jet tagging performance working point: reaching 40% efficiency and 40% purity.
- ▶ Low fake high-DCA tail background
 - *b*-jets are rare (0.1%-1%) object identified via displaced vertex, therefore sensitive to rare large-DCA fake track background.
 - The working point of B-jet tagger is few-sigma above DCA peak, and
 - Possible specification: true large DCA track/fake large DCA track > 1:1-1:few for DCA tail integrated from 2-sigma to 1mm
- ▶ Tracking efficiency
 - Efficiency for multi-track tagging algorithm is sensitive to (tracking efficiency)^N
 - Possible specification: Require 60% (HFT KPP) – 75% (HFT UPP) single track efficiency $p_T > 1$ GeV/c
- ▶ DCA
 - B-jet DCA requirement is relatively moderate
 - Requirement: DCA < 100 μm @ $p_T > 4$ GeV/c (sPHENIX proposal)
- ▶ DAQ output event rate
 - Statistical limited measurement
 - B-jets are jet-structure study based on inclusive jets, require large jet-sample rate
 - Requirement: 15 kHz trigger rate to match sPHENIX DAQ



b-jet tagging – High DCA track counting

Update in HIJING embedding:


- ▶ Haiwang Yu (NMSU) initiated the study by embedding pythia-8 MB jets into 0-4 fm HIJING background, then go through full tracking Geant4 simulation and reconstruction
- ▶ Two configuration in study
 - Cylindrically modelled MAPS + INTT + TPC (target configuration)
 - 7-layers of MAPS (Same MAPS inner tracker + MAPS outer tracker) as reference of ultimate tracking configuration of very low fake tracking rate but same physics background.




MAPS + INTT + TPC, $p_T > 1$ GeV/c

DCA_2D (cm)

Region of analysis




7-layer MAPS, $p_T > 1$ GeV/c




b-jet simulations, drawing to Geant4

sPHENIX Geant4 simulation of $p_T=30$ GeV/c B^+ -hadron





Design to Simulation



Luminosity counting

- ▶ Current RAA plot assumed 200B MB Au+Au in $|z| < 10\text{cm}$
 - 100B MB Au+Au in $|z| < 10\text{cm}$ assumed for sPHENIX proposal
 - 200B MB Au+Au in $|z| < 10\text{cm}$ following updated CAD projection
 - Will follow the final luminosity number determined by collaboration for QM17 - Gunther
- ▶ For MAPS proposal, we need updated model R_{AA} for RHIC energy




Simulation resources

- ▶ Currently we separate jet and b-tagging simulations to help speed up simulation. Need to verify factorization in the next stage
- ▶ Simulation setup used in analysis:
 - Tracking simulation in p+p in MAPS+IT+TPC (few minute / event)
 - Tracking simulation in HIJING + embedding for 7-layer MAPS (few minute / event, used for initial tunings) and for MAPS+IT+TPC (1-hour / event, use for performance plots)
- ▶ In developments
 - Silicon detectors in ladder geometry <- make available soon?
 - Pile up simulation <- make available soon?
 - TPC distortion corrections



Jet flavor definition tools

- ▶ **Unifying truth definition** and jet sample generations
 - Based on Dennis' work defining a truth tagging module run on MB events to synchronize *B*-jet definition and yield between analyzers
 - Two options provided in defining truth jet by matching *b*-quark in jet (CMS definition) or by matching *B*-hadron in jet (proposal definition)
 - Available on GitHub:
<https://github.com/sPHENIX-Collaboration/analysis/tree/master/HF-Jet/TruthGeneration>
- ▶ In collaboration with TS TG: Plan to be generalized to light-parton tagging and parton interaction channel categorizations
- ▶ **Mid-term goal**: cross checked with data and NLO generators




Jet finding and fake rejections

- ▶ HF-jet are based on jet, relying on jet finding development lead by JS TG
 - Emphasis on purity and reach to lowest-possible- p_T jet, where mass effect is maximized
 - No statistics for b -jet beyond $p_T > 50$ GeV/c
- ▶ HF-jet specific: response in detector for b -favored jet, unfolding and media modification
 - Require join study with JS TG in term of experience and toolkit developments



HF-jet TG high priority longer-term tasks

- ▶ Goal: realistic study of HF jet performance in sPHENIX simulation and reconstruction.
- ▶ High priority development tasks :
(current developers and **your help/ideas welcomed!**)
 - Realistic implementation in Geant4
 - Tony F./Gaku M./Chris P.: merged to main repository last week. Validating for general use.
 - Generalized Kalman filter
 - Haiwang Y./Chris P., ready, used in analysis, improving details
 - Multi-vertexing/ b -tagging via secondary vertexing in jet
 - Sanghoon L./Haiwang Y.: ready, used in analysis, push towards HI analysis
 - b -jet tagging: Track Counting
 - Haiwang Y./Dennis P.: ready, used in analysis, push towards 3-D DCA and HI analysis
 - b -jet tagging: Soft Lepton Tagging, exploratory
 - b -quark jet selection: B -Meson Tagging. Exploratory, volunteers from LANL and LBNL
- ▶ Area of overlapping with to other TG groups
 - JS TG: Jet detection / modern jet structure tools / event and jet flavor tagger
 - Quarkonia TG: tracking development/ HF-meson detection



Cross section from pythia8

